

Lecture #7: Particle Motion in Temporally Varying \underline{B} Fields & Adiabatic Invariance Howes ①

I Particle Motion in a Temporally Varying Magnetic Field $\underline{B}(t)$


A. Uniform Magnetic field changing in time $\underline{B}(t) = B(t) \hat{z}$.

1. Unlike the static case, Faraday's Law tells us


$$\frac{\partial \underline{B}}{\partial t} = -\nabla \times \underline{E}, \text{ so an electric field is produced.}$$

2. What will this Electric field \underline{E} do?

a. Take $\frac{dB}{dt} > 0$



b. We expect the electric field can accelerate ions or electrons.



c. For $\frac{dB}{dt} > 0$, ions are accelerated in $-\hat{\phi}$ direction
 electrons are accelerated in $+\hat{\phi}$ direction } Both gain energy.

3. Lorentz Force Law: $m \frac{d\underline{v}}{dt} = q (\underline{E} + \underline{v} \times \underline{B})$

a. Take $\underline{v} \cdot \underline{v} = v^2 \Rightarrow m \underline{v} \cdot \frac{d\underline{v}}{dt} = q \underline{v} \cdot (\underline{E} + \underline{v} \times \underline{B})$

$$\frac{d}{dt} \left(\frac{1}{2} m v^2 \right) = q \underline{v} \cdot \underline{E}$$

b. For $\frac{dB}{dt} = \frac{dB}{dt} \hat{z} = \left(\frac{\partial E_y}{\partial x} - \frac{\partial E_x}{\partial y} \right) \hat{z} \Rightarrow E_z = 0$

c. Therefore $\frac{d}{dt} \left(\frac{1}{2} m v^2 \right) = \frac{d}{dt} \left(\frac{1}{2} m v_{\perp}^2 \right)$ since $\frac{dv_{\parallel}}{dt} = 0$.

f. What is the energy change due to $\frac{dB}{dt} \neq 0$?

a. $\frac{d}{dt} \left(\frac{1}{2} m v_{\perp}^2 \right) = q \underline{v} \cdot \underline{E}$

b. If $\frac{dB}{dt}$ changes slowly, we can calculate this energy change along the unperturbed Larmor orbit.


$$\int_0^{2\pi} \frac{d}{dt} \left(\frac{1}{2} m v_{\perp}^2 \right) dt = q \int_0^{2\pi} \underline{E} \cdot \underline{v} dt = q \int_0^{2\pi} \underline{E} \cdot \frac{d\underline{r}}{dt} dt$$

Lecture #7 (Continued)
 IA.4, (Continued)

Howes 2


c. $\Delta\left(\frac{1}{2}mv_{\perp}^2\right) = q \oint_0^{2\pi/\omega_c} \underline{E} \cdot \frac{d\underline{l}}{dt} dt = q \oint_C \underline{E} \cdot d\underline{l}$

↑
Change over 1 orbit



Line integral over the path of the particle

d. By Stokes's Theorem, $q \oint_C \underline{E} \cdot d\underline{l} = q \int_S \nabla \times \underline{E} \cdot d\underline{A} = -q \int_S \frac{d\underline{B}}{dt} \cdot d\underline{A}$



Surface integral over area enclosed by the Larmor orbit

e. NOTE: For the ion motion above, $d\underline{A} = -dA \hat{z}$ (right-hand rule), so

$$\Delta\left(\frac{1}{2}mv_{\perp}^2\right) = -q \int_S \frac{d\underline{B}}{dt} \cdot \hat{z} \cdot (-dA \hat{z}) = +q \int_S \frac{dB}{dt} dA = q \frac{dB}{dt} (\pi r_L^2)$$

f. If we assume the rate of energy change is approximately constant over Larmor orbit, $\Delta\left(\frac{1}{2}mv_{\perp}^2\right) = \frac{dW_{\perp}}{dt} \Delta t = \frac{dW_{\perp}}{dt} \left(\frac{2\pi}{\omega_c}\right)$
 where $W_{\perp} = \frac{1}{2}mv_{\perp}^2$ is perpendicular energy.

g. Thus

$$\frac{dW_{\perp}}{dt} = \frac{2\omega_c}{2\pi} q \frac{dB}{dt} \pi \left(\frac{v_{\perp}^2}{\omega_c^2}\right) = \frac{q v_{\perp}^2 m}{2 e B} \frac{dB}{dt} = \left(\frac{mv_{\perp}^2}{2B}\right) \frac{dB}{dt} = \mu \frac{dB}{dt}$$

← $v_{\perp}^2 = \frac{v^2}{\omega_c^2}$

h. Since $\mu = \frac{W_{\perp}}{B}$, ~~we have~~ $\frac{dW_{\perp}}{dt} = \frac{W_{\perp}}{B} \frac{dB}{dt} \Rightarrow \left(\frac{dW_{\perp}}{W_{\perp}} - \frac{dB}{B}\right) = 0$

$$\Rightarrow \ln W_{\perp} = \ln B + C \Rightarrow \frac{W_{\perp}}{B} = \text{constant}$$

5. Therefore, for slowly varying magnetic fields $\underline{B}(t)$,


$\frac{dW_{\perp}}{dt} = 0$

I. (Continued)

B. Magnetic Flux Interpretation

1. Conservation of the Magnetic Moment μ is equivalent to maintaining a constant magnetic flux through Larmor orbits.


a. $\Phi_B = \int dA \cdot \underline{B}$



For an case $\Phi_B = B \Delta A = B \pi r^2 = \frac{\pi v_{\perp}^2}{\omega c^2} B = \pi \frac{v_{\perp}^2 m^2}{q^2 B^2} B = \frac{2\pi m}{q^2} \left(\frac{mv_{\perp}^2}{2B} \right)$

$$\Phi_B = \frac{2\pi m}{q^2} \mu$$

2. This holds for a \underline{B} -field varying (slowly) in either time or space.



$$\Phi_{B1} = \Phi_{B2}$$

II. Adiabatic Invariance

A. General Result from Hamiltonian Mechanics

For "nearly" periodic system
& slowly varying parameters,

Action Integral

$$J = \oint p dq$$

is an adiabatic invariant

i. p & q are conjugate
momentum & position coordinates.


II. (Continued)

B. Example: Harmonic Oscillator

1. Consider a time dependent harmonic oscillator.

$$\frac{d^2 x}{dt^2} + \omega^2(t) x = 0$$

Ex: Spring-mass system



$$\omega^2 = \frac{k}{m}$$

2. Position $q = x = A \sin \omega t$

Momentum $p = mv_x = m \omega A \cos \omega t$

3. Action Integral:

$$J = \oint p dq = \int_0^{2\pi/\omega} m \omega A \cos \omega t d(A \sin \omega t) = m A^2 \omega^2 \int_0^{2\pi/\omega} \cos^2 \omega t dt$$

$$= m A^2 \omega^2 \frac{\pi}{\omega} = \pi m \omega A^2$$

a. Thus $J = \pi m \omega A^2 = \text{constant}$ if $\omega(t)$ changes slowly.

b. So amplitude $A \propto \omega^{-1/2}$. If frequency decreases, amplitude will increase.

4. Total Energy $W = \frac{p^2}{2m} = \frac{1}{2} m \omega^2 A^2$, so this can also be written

$$J = 2\pi \frac{W}{\omega} = \text{constant.}$$

C. How slow must system change to satisfy invariance?

1. Since amplitude $A \propto \frac{1}{\omega^{1/2}}$, consider the WKB solution

$$X_{\text{WKB}} = \frac{1}{\sqrt{\omega(t)}} e^{\pm i \int^t \omega(t') dt'}$$

a. In this case, $J = \pi m \omega A^2$ is precisely constant.

2. This solution is an exact solution of the differential equation

$$\frac{d^2 X_{\text{WKB}}}{dt^2} + \left[\omega^2 + \frac{\ddot{\omega}}{2\omega} - \frac{3}{4} \left(\frac{\dot{\omega}}{\omega} \right)^2 \right] X_{\text{WKB}} = 0$$

Lecture 7 (Continued)

II. C2 (Continued)

b. Here $\dot{\omega} = \frac{d\omega}{dt}$ and $\ddot{\omega} = \frac{d^2\omega}{dt^2}$.

3. The WKB solution is ~~valid~~ a good approximation when

$$\omega^2 \gg \left| \frac{3}{4} \left(\frac{\dot{\omega}}{\omega} \right)^2 - \frac{\ddot{\omega}}{2\omega} \right|$$


Rule of thumb

The adiabatic invariant is approximately constant when the change of characteristic frequency is small over one period.


D. Example: Magnetic Mirror and its first, second, and third ^{adiabatic} invariants

1. Three types of periodic motion in an axisymmetric magnetic mirror

a. Larmor Motion




b. Parallel Bounce Motion



c. Azimuthal Drift Motion

(due to ∇B and Curvature Drifts)



2. First Adiabatic Invariant:

a. As we know from lecture #3, the lowest order motion in a magnetic field is Larmor motion,

$$\frac{d^2x}{dt^2} = -\omega_c^2 x \quad \text{or} \quad \frac{d^2x}{dt^2} = -\omega_c^2 x$$

Lecture #7 (Continued)
 II: D, 1 (Continued)

Hines 6

b. In this case, the action integral is

$$J_1 = \pi m \omega_c v_L^2$$

using $x = r_L \sin \omega_c t$
 $v_x = r_L \omega_c \cos \omega_c t$

c. Note that this can be written

$$J = \pi m \frac{qB}{m} \frac{v_L^2}{\left(\frac{qB}{m}\right)^2} = \frac{2\pi m}{q} \left(\frac{m v_L^2}{2B}\right) = \frac{2\pi m}{q} \mu$$

This is just the same as μ (with a constant factor)

3. Second Adiabatic Invariant, (Parallel Bounce Motion)

a. The action integral for parallel bounce motion

$$J = m \oint v_{||} ds \quad s \equiv \text{distance along magnetic field.}$$

b. We know, for a turning point at $B = B_t$,

$$\frac{1}{2} m v_{||}^2 + \mu B(s) = \mu B_t \quad (\text{Lecture #6})$$

so $v_{||}(s) = \pm \sqrt{\frac{2\mu}{m} (B_t - B(s))}$


c. This gives

$$J_2 = \sqrt{2\mu m} \oint \sqrt{B_t - B(s)} ds$$

d. Thus, for a given magnetic field configuration with $B(s)$,

$$\int_{s_a}^{s_b} \sqrt{B_t - B(s)} ds = \text{constant}$$

For bounce motion between two points s_a & s_b




II. D3C (Continued)

e. As illustrated above, the constancy of J_z (for slowly varying system parameters) can be used to determine new motion of a system.

- i. For an initial magnetic field $B_i(s)$ and initial energy, we may calculate J_z and S_{ai} & S_{bi} .
- ii. Let the magnetic field change (slowly) from $B_i(s)$ to $B_f(s)$.
- iii. Since $J_z = \int_{S_{ai}}^{S_{bi}} \sqrt{B_{fi} - B(s)} ds = \int_{S_{af}}^{S_{bf}} \sqrt{B_{ff} - B(s)} ds$, we can adjust B_{ff} (and find corresponding mirror points S_{af} & S_{bf}) until this integral is satisfied using $B_f(s)$.
- iv. The final total energy is then μB_{ff} (since $\mu = \text{const}$).

4. Third Adiabatic Invariant, (Azimuthal Drift Motion)


a. This invariant only exists in axially symmetric cases, such that the drift orbits ~~else~~ ^{of the guiding centers} are nearly closed.



b. What happens when $B(t)$ changes in time?

i. $\int_C \mathbf{E} \cdot d\mathbf{l} = - \int_S \frac{\partial B}{\partial t} \cdot d\mathbf{A}$ is change in energy

ii. Assuming Axisymmetry, $E(2\pi R) = - \frac{dB}{dt} \pi R^2 \Rightarrow E = - \frac{R}{2} \frac{dB}{dt}$



Lecture #7 (Continued)

Haves (8)

II. D. 4. b. (Continued)

$$iv. \quad \underline{v}_E = \frac{\underline{E} \times \underline{B}}{B^2} = \frac{-R \frac{dB}{dt} \hat{\phi} \times B_r \hat{z}}{B^2} = -\frac{R}{2B} \frac{dB}{dt} \hat{r}$$

$$B_r \underline{v}_E = \frac{dR}{dt} \Rightarrow \frac{dR}{dt} = -\frac{R}{2B} \frac{dB}{dt} \Rightarrow \frac{2dR}{R} = -\frac{dB}{B}$$

v. Thus $R^2 B = \text{const.}$

vi. The Magnetic Flux through drift orbit is

$$\Phi_B = \pi R^2 B = \text{const.}$$


(Assuming B is relatively constant near axis of symmetry.)

vii. Thus, the 3rd Adiabatic Invariant means the

Flux enclosed by drift orbit remains constant

Particle remains on the surface of a flux tube.

E. Example: Magnetosphere



1. Second Adiabatic Invariant applies even without axisymmetry.
2. a. For a constant Magnetic field ($\frac{dB}{dt} = 0$), energy is conserved.


$$E = \frac{1}{2} m v_{||}^2 + \mu B(s)$$

b. Since $E = \text{const}$, $B_r = \text{constant}$.

c. Following arc B_r , $I = \int_a^b \sqrt{E - \frac{\mu(s)}{B_r}} ds = \text{const.}$

3. Higher order multiple resonances

Quasi-spherical surface where particles mirror



a. Constant B_r & I mean that drifting particles remain on a surface \Rightarrow L-shell