
29:006—EXERCISES ON SIMPLE HARMONIC MOTION


Formulas

- Kinetic Energy $KE = \frac{1}{2} mv^2$
- gravitational potential energy $GPE = mgh$
- frequency $f = 1/T$, where T is the period of oscillation
- Period of a mass-spring oscillator: $T = 2\pi\sqrt{\frac{m}{k}}$
- period of a pendulum: $T = 2\pi\sqrt{\frac{L}{g}}$

Exercises

1. A 10 kg mass is attached to a rope that is 100 m long to form a huge pendulum. The mass is pulled aside so that it is 5 meters above its resting point.

- How much potential energy (PE) does it have when it is 5 m above its resting point?
- When the pendulum is released how much kinetic energy (KE) will it have when it passes through its lowest point? *Hint*: energy is conserved.
- How fast will it be moving when it passes through its lowest point?
- If it takes 2 seconds for the pendulum to reach its lowest point after it is released, when will it return to its initial position? What is this time called?



- The period, T of a harmonic oscillator is 4 seconds. What is its frequency f ?
- The frequency, f of a harmonic oscillator is 0.1 Hz. What is its period of oscillation?
- What force is needed to keep a spring stretched by 10 cm if the spring constant is 20 N/m?

5. When cart of mass m is connected to a hoop spring of spring constant k on the air track, the cart undergoes simple harmonic motion with a period of 5 seconds. The experiment is repeated with a different cart of mass M and it is found that the period is 10 seconds. What is the relationship between m and M ?
6. A huge pendulum is made by hanging a 100 kg mass at the end of a rope that is 40 m long.
- What is the period of this pendulum?
 - How many *complete* cycles will this pendulum execute in one minute?
-

Solutions

1. (a) $PE = m g h = 10 \text{ kg} \cdot 10 \text{ m/s}^2 \cdot 5 \text{ m} = 500 \text{ J}$

(b) all of the PE is converted to KE at the bottom, so $KE = PE = 500 \text{ J}$

(c) $KE = \frac{1}{2} m v^2 \rightarrow v = \sqrt{\frac{2 \cdot KE}{m}} = \sqrt{\frac{2 \cdot 500 \text{ J}}{10 \text{ kg}}} = \sqrt{\frac{1000}{10}} = \sqrt{100} = 10 \text{ m/s}$

(d) It takes 2 seconds to get to the bottom, another 2 seconds to rise to its highest point on the left side, 2 seconds to get back down to the bottom and another 2 seconds to get back to its starting point, so the total is $2 + 2 + 2 + 2 = 8$ seconds. This time is called the period, T of oscillation.

2. $f = \frac{1}{T} = \frac{1}{4 \text{ s}} = \frac{1}{4} \text{ Hz} = 0.25 \text{ Hz}$

3. $T = \frac{1}{f} = \frac{1}{0.1 \text{ Hz}} = \frac{1}{\frac{1}{10} \text{ Hz}} = 10 \text{ s}$

4. Magnitude of force, $F = k(\text{N/m}) \cdot x$ (amount of stretch in m) = $20 \text{ N/m} \cdot 0.10 \text{ m} = 2 \text{ N}$.

5. $T = 2\pi \sqrt{\frac{m}{k}} \rightarrow$ to double T , the mass must increase by a factor of 4, since $\sqrt{4} = 2$.

Therefore $M = 4 \cdot m$.

6. (a) $T = 2\pi \sqrt{\frac{L}{g}} = 2\pi \sqrt{\frac{40 \text{ m}}{10 \text{ m/s}^2}} = 2\pi \sqrt{4} = 4\pi = 12.6 \text{ s}$

(b) $f = 1/T = 0.079 \text{ Hz}$ or 0.079 cycles per second, thus in one minute (60 s) this pendulum will execute $0.079 \text{ cycles/s} \times 60 \text{ s} = 4.76 \text{ cycles}$, or 4 complete cycles